

Estimated Annual Impact Of Indigenous Foodborne Disease
- By Pathogen, England And Wales

PATHOGEN	CASES	DEATHS
<i>Campylobacter</i> spp.	337,655	80
<i>Clostridium perfringens</i>	168,436	177
<i>Yersinia</i> spp.	129,338	3
Salmonella (non-typhoidal)	73,193	209
Non-VTEC <i>E. coli</i>	62,050	6
Norovirus	61,584	10
<i>Bacillus cereus</i>	10,717	0
<i>Staphylococcus aureus</i>	9,196	0
<i>Giardia lamblia</i>	1,999	0
<i>Cryptosporidium parvum</i>	1,699	3
<i>Cyclospora cayetanensis</i>	1,026	0
<i>E. coli</i> O157:H7	1,026	23
<i>Shigella</i> spp.	308	0
Other Vibrio	291	2
<i>Listeria monocytogenes</i>	221	78
<i>Vibrio cholerae</i>	194	0
Non-O157:H7 EHEC (Enterohaemorrhagic <i>E. coli</i>)	114	3
Salmonella Paratyphi	91	0
Salmonella Typhi	86	0

Source: Disease Risks from Foods, England & Wales, 1996 - 2000, G K Adak et. al, Emerging Infectious Diseases, Vol. 11, No. 3, March 2005